


**1** Two particles  $P$  and  $Q$  of masses  $0.2\text{ kg}$  and  $0.3\text{ kg}$  respectively are free to move in a horizontal straight line on a smooth horizontal plane.  $P$  is projected towards  $Q$  with speed  $0.5\text{ m s}^{-1}$ . At the same instant  $Q$  is projected towards  $P$  with speed  $1\text{ m s}^{-1}$ .  $Q$  comes to rest in the resulting collision.

Find the speed of  $P$  after the collision.

[3]

Dotted lines for student response.



2 A car of mass 1400 kg is travelling at constant speed up a straight hill inclined at  $\alpha$  to the horizontal, where  $\sin \alpha = 0.1$ . There is a constant resistance force of magnitude 600 N. The power of the car's engine is 22 500 W.

(a) Show that the speed of the car is  $11.25 \text{ m s}^{-1}$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The car, moving with speed  $11.25 \text{ m s}^{-1}$ , comes to a section of the hill which is inclined at  $2^\circ$  to the horizontal.

(b) Given that the power and resistance force do not change, find the initial acceleration of the car up this section of the hill. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....


.....

.....

.....



3



A particle  $Q$  of mass  $0.2\text{ kg}$  is held in equilibrium by two light inextensible strings  $PQ$  and  $QR$ .  $P$  is a fixed point on a vertical wall and  $R$  is a fixed point on a horizontal floor. The angles which strings  $PQ$  and  $QR$  make with the horizontal are  $60^\circ$  and  $30^\circ$  respectively (see diagram).

Find the tensions in the two strings.

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....



4



An elevator moves vertically, supported by a cable. The diagram shows a velocity-time graph which models the motion of the elevator. The graph consists of 7 straight line segments.

The elevator accelerates upwards from rest to a speed of  $2 \text{ m s}^{-1}$  over a period of 1.5 s and then travels at this speed for 4.5 s, before decelerating to rest over a period of 1 s.

The elevator then remains at rest for 6 s, before accelerating to a speed of  $V \text{ m s}^{-1}$  downwards over a period of 2 s. The elevator travels at this speed for a period of 5 s, before decelerating to rest over a period of 1.5 s.

- (a) Find the acceleration of the elevator during the first 1.5 s. [1]

.....

.....

.....

.....

- (b) Given that the elevator starts and finishes its journey on the ground floor, find  $V$ . [2]

.....

.....

.....

.....

.....


.....

.....





5



A block of mass 5 kg is being pulled along a rough horizontal floor by a force of magnitude  $X$  N acting at  $30^\circ$  above the horizontal (see diagram). The block starts from rest and travels 2 m in the first 5 s of its motion.

- (a) Find the acceleration of the block. [2]

.....

.....

.....

.....

.....

.....

.....

- (b) Given that the coefficient of friction between the block and the floor is 0.4, find  $X$ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



.....

.....

.....

.....

.....

The block is now placed on a part of the floor where the coefficient of friction between the block and the floor has a different value. The value of  $X$  is changed to 25, and the block is now in limiting equilibrium.

- (c) Find the value of the coefficient of friction between the block and this part of the floor. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



6 A particle moves in a straight line. It starts from rest from a fixed point  $O$  on the line. Its velocity at time  $t$  s after leaving  $O$  is  $v$  m s<sup>-1</sup>, where  $v = t^2 - 8t^{\frac{3}{2}} + 10t$ .

(a) Find the displacement of the particle from  $O$  when  $t = 1$ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....







7



Two particles  $P$  and  $Q$  of masses  $0.5 \text{ kg}$  and  $m \text{ kg}$  respectively are attached to the ends of a light inextensible string. The string passes over a fixed smooth pulley which is attached to the top of two inclined planes. The particles are initially at rest with  $P$  on a smooth plane inclined at  $30^\circ$  to the horizontal and  $Q$  on a plane inclined at  $45^\circ$  to the horizontal. The string is taut and the particles can move on lines of greatest slope of the two planes. A force of magnitude  $0.8 \text{ N}$  is applied to  $P$  acting down the plane, causing  $P$  to move down the plane (see diagram).

(a) It is given that  $m = 0.3$ , and that the plane on which  $Q$  rests is smooth.

Find the tension in the string.

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

