

1 Solve the inequality $|2x - 1| > 3|x + 2|$.

[4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



3 (a) Show that the equation

$$\ln(1 + e^{-x}) + 2x = 0$$

can be expressed as a quadratic equation in e^x . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence solve the equation $\ln(1 + e^{-x}) + 2x = 0$, giving your answer correct to 3 decimal places. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



4 The equation of a curve is $y = x \tan^{-1}\left(\frac{1}{2}x\right)$.

(a) Find $\frac{dy}{dx}$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) The tangent to the curve at the point where $x = 2$ meets the y -axis at the point with coordinates $(0, p)$.

Find p . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



5 By first expressing the equation

$$\tan \theta \tan(\theta + 45^\circ) = 2 \cot 2\theta$$

as a quadratic equation in $\tan \theta$, solve the equation for $0^\circ < \theta < 90^\circ$. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....


.....

.....

.....



A series of horizontal dotted lines for writing.



- 6 (a) By sketching a suitable pair of graphs, show that the equation $x^5 = 2 + x$ has exactly one real root. [2]

.....
.....

- (b) Show that if a sequence of values given by the iterative formula

$$x_{n+1} = \frac{4x_n^5 + 2}{5x_n^4 - 1}$$

converges, then it converges to the root of the equation in part (a). [2]

.....
.....
.....
.....
.....
.....
.....



.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use the iterative formula with initial value $x_1 = 1.5$ to calculate the root correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



7 Let $f(x) = \frac{2}{(2x - 1)(2x + 1)}$.

(a) Express $f(x)$ in partial fractions. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Using your answer to part (a), show that

$$(f(x))^2 = \frac{1}{(2x - 1)^2} - \frac{1}{2x - 1} + \frac{1}{2x + 1} + \frac{1}{(2x + 1)^2}. \quad [2]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....


.....

.....



(c) Hence show that $\int_1^2 (f(x))^2 dx = \frac{2}{5} + \frac{1}{2} \ln\left(\frac{5}{9}\right)$. [5]

Dotted lines for answer writing.



(b) Find angle BAD , giving your answer in degrees. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Find the area of the parallelogram correct to 3 significant figures. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....




- (b) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities

$$|z - 2 - 2i| \leq 2, \quad 0 \leq \arg z \leq \frac{1}{4}\pi \quad \text{and} \quad \operatorname{Re} z \leq 3. \quad [5]$$



10



A tank containing water is in the form of a hemisphere. The axis is vertical, the lowest point is A and the radius is r , as shown in the diagram. The depth of water at time t is h . At time $t = 0$ the tank is full and the depth of the water is r . At this instant a tap at A is opened and water begins to flow out at a rate proportional to \sqrt{h} . The tank becomes empty at time $t = 14$.

The volume of water in the tank is V when the depth is h . It is given that $V = \frac{1}{3}\pi(3rh^2 - h^3)$.

(a) Show that h and t satisfy a differential equation of the form

$$\frac{dh}{dt} = -\frac{B}{2rh^{\frac{1}{2}} - h^{\frac{3}{2}}},$$

where B is a positive constant. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



(b) Solve the differential equation and obtain an expression for t in terms of h and r . [8]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

