## Solomon Practice Paper

Core Mathematics 1A

Time allowed: 90 minutes

Centre: www.CasperYC.club

Name:

Teacher:

How	Ι	can	achieve	better:
	- C	oun	actilicite	

- •

- •

CC	•	\$	
	BY	NC	ND

July 14, 2025



Question	Points	Score
1	4	
2	4	
3	5	
4	6	
5	7	
6	8	
7	9	
8	10	
9	11	
10	11	
Total:	75	

re Mathematics – Practice Paper 1A (a) Express $\frac{21}{\sqrt{7}}$ in the form $k\sqrt{7}$ .	Page 1 of 10
(a) Express $\frac{1}{\sqrt{7}}$ in the form $k \sqrt{7}$ . (b) Express $8^{-\frac{1}{3}}$ as an exact fraction in its simplest form.	
(b) Express 8 <sup>-3</sup> as an exact fraction in its simplest form.	Total
	10000
	$\sim$
	<u>Joo</u>
	$( \circ)$

[4]

2. Evaluate

$$\sum_{r=10}^{30} 7 + 2r$$



www.CasperYC.club

 $6x^2 - 1$ 

www.CasperYC.club

4. (a) Solve the inequality x<sup>2</sup> + 3x > 10. [3]
(b) Find the set of values of x which satisfy both of the following inequalities: [3]
3x - 2 < x + 3 and x<sup>2</sup> + 3x > 10.

Total: 6



www.CasperYC.club

5. The sequence  $u_1, u_2, u_3, \ldots$  is defined by the recurrence relation

$$u_{n+1} = (u_n)^2 - 1, \quad n \ge 1.$$

Given that  $u_1 = k$ , where k is a constant,

- (a) find expressions for  $u_2$  and  $u_3$  in terms of k.
- Given also that  $u_2 + u_3 = 11$ ,
- (b) find the possible values of k.

Last updated: July 14, 2025



[3]

[4]


Total: 7

$$x^2 + 4kx - k = 0.$$

(b) Hence find the set of values of k for which the equation has no real roots. [4]

[4]

www.CasperYC.club



7. (a) Describe fully a single transformation that maps the graph of y = <sup>1</sup>/<sub>x</sub> onto the graph of y = <sup>3</sup>/<sub>x</sub>. [2]
(b) Sketch the graph of y = <sup>3</sup>/<sub>x</sub> and write down the equations of any asymptotes. [3]
(c) Find the values of the constant c for which the straight line y = c - 3x is a tangent to the curve y = <sup>3</sup>/<sub>x</sub>.

Total: 9



www.CasperYC.club


## Core Mathematics – Practice Paper 1A

8. The points $P$ and $Q$ have coordinates $(7, 4)$ and $(9, 7)$ respectively.	
(a) Find an equation for the straight line $l$ which passes through $P$ and $Q$ . Give your answer in the form $ax + by + c = 0$ , where $a, b$ and $c$ are integers.	[4]
The straight line $m$ has gradient 8 and passes through the origin, $O$ .	
(b) Write down an equation for $m$ .	[1]
The lines $l$ and $m$ intersect at the point $R$ .	
(c) Show that $OP = OR$ .	[5]
	Total: 10



www.CasperYC.club

9. Figure below shows the curve with equation y = f(x) which crosses the x-axis at the origin and at the points A and B.



Given that

$$f'(x) = 6 - 4x - 3x^2,$$

- (a) find an expression for y in terms of x,
- (b) show that  $AB = k\sqrt{7}$ , where k is an integer to be found.

[5]

[6]

Total:	11

www.CasperYC.club

10. A curve has the equation  $y = x + \frac{3}{x}, x \neq 0$ .

The point P on the curve has x-coordinate 1.

- (a) Show that the gradient of the curve at P is -2.
- (b) Find an equation for the normal to the curve at P, giving your answer in the form y = mx + c. [4]
- (c) Find the coordinates of the point where the normal to the curve at P intersects the curve [4] again.

Total: 11

[3]

www.CasperYC.club

