

Solomon Practice Paper

Core Mathematics 1H

Time allowed: 90 minutes

Centre: www.CasperYC.club

Name:

Teacher:

Question	Points	Score
1	3	
2	4	
3	6	
4	7	
5	7	
6	8	
7	8	
8	9	
9	10	
10	13	
Total:	75	

How I can achieve better:

-
-
-




Last updated: July 14, 2025



1. Evaluate


[3]

$$\sum_{r=1}^{30} (3r + 4).$$




2. (a) Express $x^2 + 6x + 7$ in the form $(x + a)^2 + b$. [3]
- (b) State the coordinates of the minimum point of the curve $y = x^2 + 6x + 7$. [1]

Total: 4



3. The straight line l_1 has the equation $3x - y = 0$.
The straight line l_2 has the equation $x + 2y - 4 = 0$.
- (a) Sketch l_1 and l_2 on the same diagram, showing the coordinates of any points where each line meets the coordinate axes. [3]
- (b) Find, as exact fractions, the coordinates of the point where l_1 and l_2 intersect. [3]


Total: 6



4. Find the pairs of values (x, y) which satisfy the simultaneous equations

[7]

$$\begin{cases} 3x^2 + y^2 &= 21 \\ 5x + y &= 7 \end{cases}$$



5. (a) Sketch on the same diagram the graphs of $y = (x - 1)^2(x - 5)$ and $y = 8 - 2x$. [5]
Label on your diagram the coordinates of any points where each graph meets the coordinate axes.

(b) Explain how your diagram shows that there is only one solution, α , to the equation [1]

$$(x - 1)^2(x - 5) = 8 - 2x.$$

(c) State the integer, n , such that [1]

$$n < \alpha < n + 1.$$

Total: 7



6. The curve with equation $y = x^2 + 2x$ passes through the origin, O .
- (a) Find an equation for the normal to the curve at O . [5]
- (b) Find the coordinates of the point where the normal to the curve at O intersects the curve again. [3]
- again.

Total: 8



7. Given that

$$y = \sqrt{x} - \frac{4}{\sqrt{x}},$$

- (a) find $\frac{dy}{dx}$, [3]
- (b) find $\frac{d^2y}{dx^2}$, [2]
- (c) show that [3]

$$4x^2 \frac{d^2y}{dx^2} + 4x \frac{dy}{dx} - y = 0.$$

Total: 8




8. (a) Prove that the sum of the first n positive integers is given by [4]

$$\frac{1}{2}n(n + 1).$$

(b) Hence, find the sum of [5]

- i. the integers from 100 to 200 inclusive,
- ii. the integers between 300 to 600 inclusive which are divisible by 3.

Total: 9




9. (a) Express each of the following in the form $p + q\sqrt{2}$ where p and q are rational. [5]
- i. $(4 - 3\sqrt{2})^2$

ii. $\frac{1}{2+\sqrt{2}}$
- (b)


i. Solve the equation $y^2 + 8 = 9y$. [5]

ii. Hence solve the equation $x^3 + 8 = 9x^{\frac{3}{2}}$.

Total: 10



10. Figure shows the curve with equation $y = f(x)$.



The curve meets the x -axis at the origin and at the point A . Given that

$$f'(x) = 3x^{\frac{1}{2}} - 4x^{-\frac{1}{2}},$$

- (a) find $f(x)$, [5]
- (b) find the coordinate of A . [2]

The point B on the curve has x -coordinate 2.

- (c) Find an equation for the tangent to the curve at B in the form $y = mx + c$. [6]

Total: 13

