

Solomon Practice Paper

Core Mathematics 2E

Time allowed: 90 minutes

Centre: www.CasperYC.club

Name:

Teacher:

Question	Points	Score
1	4	
2	5	
3	6	
4	8	
5	9	
6	9	
7	10	
8	12	
9	12	
Total:	75	

How I can achieve better:

-
-
-



Last updated:

July 14, 2025



1. Evaluate

[4]


$$\int_2^4 2 - \frac{1}{x^2} \, dx.$$



[5]

Find the set of values of x for which $f(x)$ is increasing.

Find the set of values of x for which $f(x)$ is increasing.




3. Given that $p = \log_2(3)$ and $q = \log_2(5)$, find expressions in terms of p and q for

(a) $\log_2(45)$. [3]

(b) $\log_2(0.3)$. [3]

Total: 6



(a) Find the value of k .

[3]


(b) show that the coefficient of x^3 in the expansion is 4375,

[2]

[3]

$$(2 - x)(1 + kx)^7.$$

Total: 8




- The finite region R is bounded by the curve $y = \cos^2(x)$, where x is measured in radians, the positive coordinate axes and the line $x = \frac{\pi}{3}$.

- The finite region S is bounded by the curve $y = \sin^2(x)$, where x is measured in radians, the positive coordinate axes and the line $x = \pi/3$.

- Total: 9



6. Figure shows triangle ABC in which $AC = 8$ cm and $\angle BAC = \angle BCA = 30^\circ$.



(a) Find the area of triangle ABC in the form $k\sqrt{3}$. [5]

The point M is the mid-point of AC and the points N and O lie on AB and BC such that MN and MO are arcs of circles with centres A and C respectively.

(b) Show that the area of the shaded region $BNMO$ is $\frac{8}{3} (2\sqrt{3} - \pi)$ cm². [4]

Total: 9



$$x^2 + y^2 + 10x - 8y + k = 0,$$

Given that the point with coordinates $(-6, 5)$ lies on C ,

- A straight line which passes through the point $A(2, 3)$ is a tangent to C at the point B .

- (c) Find the length AB in the form $k\sqrt{3}$. [5]

Total: 10



For this scheme,

- (c) Find, to the nearest pound, how much more or less will be in the account at the end of the eighth year under this scheme. [5]

Total: 12



$$f(x) = x^3 + kx^2 - 7x - 15,$$

When $f(x)$ is divided by $(x + 1)$ the remainder is r .

When $f(x)$ is divided by $(x - 3)$ the remainder is $3r$.

- Find the value of k . [5]
- Find the value of r . [1]
- Show that $(x - 5)$ is a factor of $f(x)$. [2]
- Show that there is only one real solution to the equation $f(x) = 0$. [4]

Total: 12

