

# Solomon Practice Paper

## Core Mathematics 4F

Time allowed: 90 minutes

Centre: [www.CasperYC.club](http://www.CasperYC.club)

Name:

Teacher:

Question	Points	Score
1	8	
2	8	
3	9	
4	9	
5	11	
6	13	
7	17	
Total:	75	

How I can achieve better:

- 
- 
- 




Last updated:

July 14, 2025



[8]

Find the coordinates of the points where the tangent to the curve is parallel to the  $x$ -axis.



2. Use the substitution  $x = 2 \tan(u)$  to show that

[8]

$$\int_0^2 \frac{x^2}{x^2 + 4} \, dx = \frac{1}{2}(4 - \pi).$$



3. (a) Show that [2]

$$\left(1\frac{1}{24}\right)^{-\frac{1}{2}} = k\sqrt{6}$$

where  $k$  is rational.


(b) Expand [4]

$$\left(1 + \frac{1}{2}x\right)^{-\frac{1}{2}}, \quad |x| < 2,$$

in ascending powers of  $x$  up to and including the term in  $x^3$ , simplifying each coefficient.

(c) Use your answer to part (b) with  $x = \frac{1}{12}$  to find an approximate value for  $\sqrt{6}$ , giving your [3]  
answer to 5 decimal places.

Total: 9




$$\mathbf{r} = (7\mathbf{j} - 4\mathbf{k}) + s(4\mathbf{i} - 3\mathbf{j} + \mathbf{k}), \quad \text{and} \quad \mathbf{r} = (-7\mathbf{i} + \mathbf{j} + 8\mathbf{k}) + t(-3\mathbf{i} + 2\mathbf{k}),$$

(a) Show that the two lines intersect and find the position vector of the point where they meet. [5]

(b) Find, in degrees to 1 decimal place, the acute angle between the lines. [4]

Total: 9



5. A curve has parametric equations

$$x = \frac{t}{2-t}, \quad \text{and} \quad y = \frac{1}{1+t}, \quad -1 < t < 2.$$

(a) Show that


$$\frac{dy}{dx} = -\frac{1}{2} \left( \frac{2-t}{1+t} \right)^2.$$

(b) Find an equation for the normal to the curve at the point where  $t = 1$ .

(c) Show that the cartesian equation of the curve can be written in the form

$$y = \frac{1+x}{1+3x}.$$

Total: 11





6. (a) Find


[3]

$$\int \tan^2(x) \, dx.$$

(b) Show that

[4]

$$\int \tan(x) \, dx = \ln |\sec(x)| + c,$$

where  $c$  is an arbitrary constant.Figure shows part of the curve with equation  $y = x^{\frac{1}{2}} \tan(x)$ .


The shaded region bounded by the curve, the  $x$ -axis and the line  $x = \frac{\pi}{3}$  is rotated through  $2\pi$  radians about the  $x$ -axis.

(c) Show that the volume of the solid formed is

[6]

$$\frac{1}{18}\pi^2 (6\sqrt{3} - \pi) - \pi \ln(2).$$


Total: 13







7. Figure shows a hemispherical bowl of radius 5 cm.



The bowl is filled with water but the water leaks from a hole at the base of the bowl. At time  $t$  minutes, the depth of water is  $h$  cm and the volume of water in the bowl is  $V$  cm<sup>3</sup>, where

$$V = \frac{1}{3}\pi h^2(15 - h).$$

In a model it is assumed that the rate at which the volume of water in the bowl decreases is proportional to  $V$ .

(a) Show that

$$\frac{dh}{dt} = -\frac{kh(15 - h)}{3(10 - h)},$$

where  $k$  is a positive constant.

(b) Express

$$\frac{3(10 - h)}{h(15 - h)}$$

in partial fractions.

Given that when  $t = 0$ ,  $h = 5$ ,


(c) show that

$$h^2(15 - h) = 250e^{-kt}.$$

Given also that when  $t = 2$ ,  $h = 4$ ,

(d) find the value of  $k$  to 3 significant figures.

Total: 17



Lined area for writing answers, consisting of 20 horizontal lines.

