

# Solomon Practice Paper

## Core Mathematics 1I

Time allowed: 90 minutes

Centre: [www.CasperYC.club](http://www.CasperYC.club)

Name:

Teacher:

Question	Points	Score
1	3	
2	3	
3	4	
4	6	
5	7	
6	8	
7	8	
8	10	
9	13	
10	13	
Total:	75	

How I can achieve better:

- 
- 
- 



Last updated: May 5, 2023



1. The  $n$ th term of a sequence is defined by [3]

$$u_n = n^2 - 6n + 11, n \geq 1.$$

Given that the  $k$ th term of the sequence is 38, find the value of  $k$ .

2. Find [3]

$$\int 4x^2 - \sqrt{x} \, dx.$$

3. Find the integer  $n$  such that [4]

$$4\sqrt{12} - \sqrt{75} = \sqrt{n}.$$

4. (a) Evaluate  $\left(36^{\frac{1}{2}} + 16^{\frac{1}{4}}\right)$ . [3]

- (b) Solve the equation  $3x^{-\frac{1}{2}} - 4 = 0$ . [3]

Total: 6

5. The curve  $y = f(x)$  passes through the point  $P(-1, 3)$  and is such that

$$\frac{dy}{dx} = -\frac{1}{x^2}, \quad x \neq 0.$$

- (a) Using integration, find  $f(x)$ . [4]

- (b) Sketch the curve  $y = f(x)$  and write down the equations of its asymptotes. [3]

Total: 7

6.

$$f(x) = x^2 - 10x + 17.$$

- (a) Express  $f(x)$  in the form  $a(x + b)^2 + c$ . [3]

- (b) State the coordinates of the minimum point of the curve  $y = f(x)$ . [1]

- (c) Deduce the coordinates of the minimum point of each of the following curves: [4]

i.  $y = f(x) + 4$ ,

ii.  $y = f(2x)$ .

Total: 8

7. Given that the equation


$$4x^2 - kx + k - 3 = 0,$$

where  $k$  is a constant, has real roots,

- (a) show that  $k^2 - 16k + 48 \geq 0$ , [2]

- (b) find the set of possible values of  $k$ , [3]

- (c) state the smallest value of  $k$  for which the roots are equal and solve the equation when  $k$  takes this value. [3]



Total: 8

8. (a) The first and third terms of an arithmetic series are 3 and 27 respectively. [5]
- Find the common difference of the series.
  - Find the sum of the first 11 terms of the series.

- (b) Find the sum of the integers between 50 and 150 which are divisible by 8. [5]

Total: 10

9. A curve has the equation  $y = x^3 - 5x^2 + 7x$ .

- (a) Show that the curve only crosses the  $x$ -axis at one point. [4]

The point  $P$  on the curve has coordinates  $(3, 3)$ .


- (b) Find an equation for the normal to the curve at  $P$ , giving your answer in the form  $ax + by = c$ , [6]  
where  $a, b$  and  $c$  are integers.

The normal to the curve at  $P$  meets the coordinate axes at  $Q$  and  $R$ .

- (c) Show that triangle  $OQR$ , where  $O$  is the origin, has area  $28\frac{1}{8}$ . [3]

Total: 13

10. Figure shows the parallelogram  $ABCD$ .



The points  $A$  and  $B$  have coordinates  $(-1, 3)$  and  $(3, 4)$  respectively and lie on the straight line  $l_1$ .

- (a) Find an equation for  $l_1$ , giving your answer in the form  $ax + by + c = 0$ , where  $a, b$  and  $c$  [4]  
are integers.

The points  $C$  and  $D$  lie on the straight line  $l_2$  which has the equation  $x - 4y - 21 = 0$ .

- (b) Show that the distance between  $l_1$  and  $l_2$  is  $k\sqrt{17}$ , where  $k$  is an integer to be found. [7]

- (c) Find the area of parallelogram  $ABCD$ . [2]

Total: 13

