

# Solomon Practice Paper

## Core Mathematics 3F

Time allowed: 90 minutes

Centre: [www.CasperYC.club](http://www.CasperYC.club)


Name:

Teacher:


Question	Points	Score
1	6	
2	7	
3	8	
4	8	
5	10	
6	11	
7	12	
8	13	
Total:	75	

How I can achieve better:

- 
- 
- 



Last updated: May 5, 2023



1. Solve the equation

[6]

$$3 \csc(\theta^\circ) + 8 \cos(\theta^\circ) = 0$$

for  $\theta$  in the interval  $0 \leq \theta \leq 180$ , giving your answers to 1 decimal place.

2. The functions
- $f$
- and
- $g$
- are defined by

$$f: x \rightarrow 1 - ax, \quad x \in \mathbb{R},$$

$$g: x \rightarrow x^2 + 2ax + 2, \quad x \in \mathbb{R},$$

where  $a$  is a constant.

- (a) Find the range of
- $g$
- in terms of
- $a$
- .

[3]

Given that  $gf(3) = 7$ ,

- (b) find the two possible values of
- $a$
- .

[4]

Total: 7

3. (a) Solve the equation

[3]

$$\ln(3x + 1) = 2$$

giving your answer in terms of  $e$ .

- (b) Prove, by counter-example, that the statement

[5]

$$\ln(3x^2 + 5x + 3) \geq 0 \text{ for all real values of } x$$

is false.

Total: 8

4. A curve has the equation
- $x = y\sqrt{1 - 2y}$
- .

- (a) Show that

[5]

$$\frac{dy}{dx} = \frac{\sqrt{1 - 2y}}{1 - 3y}.$$

The point  $A$  on the curve has  $y$ -coordinate  $-1$ .

- (b) Show that the equation of tangent to the curve at
- $A$
- can be written in the form

[3]

$$\sqrt{3}x + py + q = 0$$


where  $p$  and  $q$  are integers to be found.

Total: 8

5. (a) Sketch the graph of

[5]

$$y = 2 + \sec\left(x - \frac{\pi}{6}\right)$$




for  $x$  in the interval  $0 \leq x \leq 2\pi$ .

Show on your sketch the coordinates of any turning points and the equations of any asymptotes.

- (b) Find, in terms of  $\pi$ , the  $x$ -coordinates of the points where the graph crosses the  $x$ -axis. [5]

Total: 10

6. Figure shows the curve  $y = f(x)$  which has a minimum point at  $(-\frac{3}{2}, 0)$ , a maximum point at  $(3, 6)$  and crosses the  $y$ -axis at  $(0, 4)$ .



Sketch each of the following graphs on separate diagrams. In each case, show the coordinates of any turning points and of any points where the graph meets the coordinate axes.

- (a)  $y = f(|x|)$  [3]  
 (b)  $y = 2 + f(x + 3)$  [4]  
 (c)  $y = \frac{1}{2}f(-x)$  [4]

Total: 11

7.

$$f(x) = 1 + \frac{4x}{2x - 5} - \frac{15}{2x^2 - 7x + 5}, x \in \mathbb{R}, x < 1.$$

- (a) Show that [5]

$$f(x) = \frac{3x + 2}{x - 1}.$$

- (b) Find an expression for the inverse function  $f^{-1}(x)$  and state its domain. [5]

- (c) Solve the equation  $f(x) = 2$ . [2]


Total: 12

8. A curve has the equation  $y = x^2 - \sqrt{4 + \ln(x)}$ .

- (a) Show that the tangent to the curve at the point where  $x = 1$  has the equation [5]

$$7x - 4y = 11.$$

The curve has a stationary point with  $x$ -coordinate  $\alpha$ .



(b) Show that  $0.3 < \alpha < 0.4$ . [3]

(c) Show that  $\alpha$  is a solution of the equation [2]

$$x = \frac{1}{2} (4 + \ln(x))^{-\frac{1}{4}}$$

(d) Use the iteration formula [3]

$$x_{n+1} = \frac{1}{2} (4 + \ln(x_n))^{-\frac{1}{4}}$$

with  $x_0 = 0.35$ , to find  $x_1, x_2, x_3$  and  $x_4$ , giving your answers to 5 decimal places.

Total: 13

