

Solomon Practice Paper

Core Mathematics 4F

Time allowed: 90 minutes

Centre: www.CasperYC.club


Name:

Teacher:


Question	Points	Score
1	8	
2	8	
3	9	
4	9	
5	11	
6	13	
7	17	
Total:	75	

How I can achieve better:

-
-
-



Last updated: May 5, 2023



1. A curve has the equation [8]

$$2x^2 + xy - y^2 + 18 = 0.$$

Find the coordinates of the points where the tangent to the curve is parallel to the x -axis.

2. Use the substitution $x = 2 \tan(u)$ to show that [8]

$$\int_0^2 \frac{x^2}{x^2 + 4} dx = \frac{1}{2}(4 - \pi).$$

3. (a) Show that [2]

$$\left(1 - \frac{1}{24}\right)^{-\frac{1}{2}} = k\sqrt{6}$$

where k is rational.

- (b) Expand [4]

$$\left(1 + \frac{1}{2}x\right)^{-\frac{1}{2}}, \quad |x| < 2,$$

in ascending powers of x up to and including the term in x^3 , simplifying each coefficient.

- (c) Use your answer to part (b) with $x = \frac{1}{12}$ to find an approximate value for $\sqrt{6}$, giving your answer to 5 decimal places. [3]

Total: 9

4. Relative to a fixed origin, two lines have the equations

$$\mathbf{r} = (7\mathbf{j} - 4\mathbf{k}) + s(4\mathbf{i} - 3\mathbf{j} + \mathbf{k}), \quad \text{and} \quad \mathbf{r} = (-7\mathbf{i} + \mathbf{j} + 8\mathbf{k}) + t(-3\mathbf{i} + 2\mathbf{k}),$$

where s and t are scalar parameters.

- (a) Show that the two lines intersect and find the position vector of the point where they meet. [5]

- (b) Find, in degrees to 1 decimal place, the acute angle between the lines. [4]

Total: 9

5. A curve has parametric equations

$$x = \frac{t}{2-t}, \quad \text{and} \quad y = \frac{1}{1+t}, \quad -1 < t < 2.$$


- (a) Show that [4]

$$\frac{dy}{dx} = -\frac{1}{2} \left(\frac{2-t}{1+t} \right)^2.$$

- (b) Find an equation for the normal to the curve at the point where $t = 1$. [3]

- (c) Show that the cartesian equation of the curve can be written in the form [4]

$$y = \frac{1+x}{1+3x}.$$



Total: 11

6. (a) Find


$$\int \tan^2(x) dx.$$

[3]

(b) Show that

$$\int \tan(x) dx = \ln |\sec(x)| + c,$$

[4]

where c is an arbitrary constant.Figure shows part of the curve with equation $y = x^{\frac{1}{2}} \tan(x)$.The shaded region bounded by the curve, the x -axis and the line $x = \frac{\pi}{3}$ is rotated through 2π radians about the x -axis.


(c) Show that the volume of the solid formed is

[6]


$$\frac{1}{18}\pi^2 (6\sqrt{3} - \pi) - \pi \ln(2).$$

Total: 13

7. Figure shows a hemispherical bowl of radius 5 cm.

The bowl is filled with water but the water leaks from a hole at the base of the bowl. At time t minutes, the depth of water is h cm and the volume of water in the bowl is V cm³, where

$$V = \frac{1}{3}\pi h^2(15 - h).$$

In a model it is assumed that the rate at which the volume of water in the bowl decreases is proportional to V .

(a) Show that

$$\frac{dh}{dt} = -\frac{kh(15-h)}{3(10-h)},$$

[5]

where k is a positive constant.

(b) Express

$$\frac{3(10-h)}{h(15-h)}$$

[3]

in partial fractions.

Given that when $t = 0$, $h = 5$,

(c) show that

$$h^2(15-h) = 250e^{-kt}.$$

[6]

Given also that when $t = 2$, $h = 4$,

(d) find the value of k to 3 significant figures.

[3]

Total: 17

