

# Solomon Practice Paper

## Core Mathematics 4L

Time allowed: 90 minutes

Centre: [www.CasperYC.club](http://www.CasperYC.club)


Name:

Teacher:


Question	Points	Score
1	8	
2	8	
3	10	
4	11	
5	12	
6	12	
7	14	
Total:	75	

How I can achieve better:

- 
- 
- 



Last updated: May 5, 2023



1. The number of people,  $n$ , in a queue at a Post Office  $t$  minutes after it opens is modelled by the differential equation

$$\frac{dn}{dt} = e^{0.5t} - 5, \quad t \geq 0.$$

- (a) Find, to the nearest second, the time when the model predicts that there will be the least number of people in the queue. [3]
- (b) Given that there are 20 people in the queue when the Post Office opens, solve the differential equation. [4]
- (c) Explain why this model would not be appropriate for large values of  $t$ . [1]

Total: 8

2. A curve has the equation

$$3x^2 + xy - 2y^2 + 25 = 0.$$

Find an equation for the normal to the curve at the point with coordinates  $(1, 4)$ , giving your answer in the form  $ax + by + c = 0$ , where  $a, b$  and  $c$  are integers.

3. (a) Use the substitution  $u = 2 - x^2$  to find [4]


$$\int \frac{x}{2 - x^2} dx.$$

- (b) Evaluate [6]

$$\int_0^{\frac{\pi}{4}} \sin(3x) \cos(x) dx.$$

Total: 10


4. Figure shows the curve with equation  $y = x\sqrt{\ln(x)}$ ,  $x \geq 1$ .



The shaded region is bounded by the curve, the  $x$ -axis and the line  $x = 3$ .

- (a) Using the trapezium rule with two intervals of equal width, estimate the area of the shaded region. [4]

The shaded region is rotated through  $360^\circ$  about the  $x$ -axis.



(b) Find the exact volume of the solid formed. [7]

Total: 11

5.

$$f(x) = \frac{5 - 8x}{(1 + 2x)(1 - x)^2}.$$

(a) Express  $f(x)$  in partial fractions. [5]


(b) Find the series expansion of  $f(x)$  in ascending powers of  $x$  up to and including the term in  $x^3$ , simplifying each coefficient. [6]

(c) State the set of values of  $x$  for which your expansion is valid. [1]

Total: 12

6. Figure shows the curve with parametric equations

$$x = t + \sin(t), \quad \text{and} \quad y = \sin(t), \quad 0 \leq t \leq \pi.$$



(a) Find  $\frac{dy}{dx}$  in terms of  $t$ . [3]

(b) Find, in exact form, the coordinates of the point where the tangent to the curve is parallel to the  $x$ -axis. [3]

(c) Show that the region bounded by the curve and the  $x$ -axis has area 2. [6]

Total: 12

7. The line  $l_1$  passes through the points  $A$  and  $B$  with position vectors  $(3\mathbf{i} + 6\mathbf{j} - 8\mathbf{k})$  and  $(8\mathbf{j} - 6\mathbf{k})$  respectively, relative to a fixed origin.

(a) Find a vector equation for  $l_1$ . [2]

The line  $l_2$  has vector equation


$$\mathbf{r} = (-2\mathbf{i} + 10\mathbf{j} + 6\mathbf{k}) + \mu(7\mathbf{i} - 4\mathbf{j} + 6\mathbf{k}),$$

where  $\mu$  is a scalar parameter.

(b) Show that lines  $l_1$  and  $l_2$  intersect. [4]

(c) Find the coordinates of the point where  $l_1$  and  $l_2$  intersect. [2]

The point  $C$  lies on  $l_2$  and is such that  $AC$  is perpendicular to  $AB$ .



(d) Find the position vector of  $C$ .

[6]

Total: 14

