

Solomon Practice Paper

Further Pure Mathematics 1C

Time allowed: 90 minutes

Centre: www.CasperYC.club

Name:

Teacher:

Question	Points	Score
1	6	
2	9	
3	9	
4	10	
5	12	
6	13	
7	16	
Total:	75	

How I can achieve better:

-
-
-



Last updated:

July 14, 2025



1. Find the set of values of x for which

[6]

$$|x - 2| > 2|x + 1|.$$




2. (a) By using the substitution $y = vx$, or otherwise, find the general solution of the differential equation [7]

$$xy \frac{dy}{dx} = x^2 + y^2.$$

- (b) Given also that $y = 2$ when $x = 1$, show that for $x > 0$ [2]

$$y^2 = 2x^2 (\ln |x| + 2).$$

Total: 9



3. (a) Find the sum of the series

[3]

$$2^3 + 4^3 + 6^3 + \dots + (2n)^3,$$

giving your answer in a simplified form.


- (b) Hence, or otherwise, show that the sum of the series

[6]

$$1^3 - 2^3 + 3^3 - 4^3 + \dots + (2n-1)^3 - (2n)^3$$

is $-n^2(4n+3)$.


Total: 9



4. Find the general solution of the differential equation

[10]

$$\frac{d^2y}{dx^2} - 6\frac{dy}{dx} + 9y = 2e^{3x}.$$



5.





Figure shows part of the curve $y = f(x)$ where

$$f(x) \equiv 2x - \tan(x), \quad x \in \mathbb{R}, \quad 0 \leq x < \frac{\pi}{2}.$$

- (a) Show that there is a root, α , of the equation $f(x) = 0$ in the interval $(1, 1.5)$. [2]
- (b) Use the Newton–Raphson method with an initial value of $x = 1.25$ to find α correct to 2 decimal places and justify the accuracy of your answer. [7]
- (c) Explain with the aid of a diagram why the Newton–Raphson method fails if an initial value of $x = 0.75$ is used when trying to find α . [3]

Total: 12



6. The complex numbers z and w are defined such that

$$\begin{cases} 3z + w = 14 \\ z - \mathbf{i}w = 15 - 9\mathbf{i} \end{cases}$$


(a) Show that $z = 3 - 4\mathbf{i}$ and find w in the form $a + \mathbf{i}b$, where a and b are real numbers.

[6]

(b) Find the square roots of z in the form $c + \mathbf{i}d$, where c and d are real numbers.

[7]

Total: 13



7.




Figure shows the curves with polar equations

$$\begin{cases} r = 4 \sin(2\theta), & 0 \leq \theta \leq \frac{\pi}{2}, \\ r = 4 \cos(\theta), & 0 \leq \theta \leq \frac{\pi}{2}. \end{cases}$$

(a) Find the polar coordinates of the point P where the two curves intersect. [5]

(b) Find the exact area of the shaded region bounded by the two curves. [11]

Total: 16

