

# Solomon Practice Paper

## Further Pure Mathematics 2A

Time allowed: 90 minutes

Centre: [www.CasperYC.club](http://www.CasperYC.club)

Name:

Teacher:

Question	Points	Score
1	5	
2	8	
3	9	
4	10	
5	10	
6	10	
7	11	
8	12	
Total:	75	

How I can achieve better:

- 
- 
- 



Last updated: July 14, 2025



1. A curve has the equation

[5]

$$y = x + 2x^2 + 5x^3.$$


Show that the radius of curvature of the curve at the origin is  $\frac{1}{\sqrt{2}}$ .



2. Show that

[8]

$$\int_0^{\ln(2)} x \operatorname{sech}^2(x) \, dx = \frac{3}{5} \ln(2) - \ln\left(\frac{5}{4}\right).$$



3. (a) Prove that

[3]

$$\frac{d}{dx} \arcsin(2x) = \frac{2}{\sqrt{1-4x^2}}.$$

Given that


$$f(x) = 2x \arcsin(2x) + \sqrt{1 - 4x^2},$$

(b) show that


[6]

$$f''(x) \left[ f(x) - xf'(x) \right] = 4.$$

Total: 9



4.



The parametric coordinates of the curve  $C$  shown are

$$x = t^2 \quad \text{and} \quad y = t - \frac{1}{3}t^3, \quad 0 \leq t \leq a.$$

The curve  $C$  meets the  $x$ -axis at the point  $A$  where  $t = a$ .

(a) Find the value of  $a$ .

[2]

The curve  $C$  is rotated through  $2\pi$  about  $Ox$ .

(b) Find the surface area of the solid generated.

[8]

Total: 10



5. (a) Using the definitions of  $\cosh x$  and  $\sinh x$  in terms of  $e^x$  and  $e^{-x}$ , prove that [3]


$$\cosh(2x) = 2 \cosh^2(x) - 1.$$

(b) Solve the equation [7]

$$2 \cosh(2x) = 13 \cosh(x) - 12,$$

giving your answers in terms of natural logarithms.

Total: 10



6.

$$x^2 - 10x + 41 \equiv (x + a)^2 + b.$$


(a) Find the values of the constants  $a$  and  $b$ . [2]

(b) Show that [8]

$$\int_5^9 \frac{x}{\sqrt{x^2 - 10x + 41}} \, dx = p \left( \sqrt{2} - 1 \right) + q \ln(r),$$

stating your values of  $p, q$  and  $r$ .

Total: 10



7.

$$I_n \int_0^{\frac{\pi}{2}} x^n \cos(x) \, dx, \quad n \geq 0.$$

(a) Prove that


[5]

$$I_n \left(\frac{\pi}{2}\right)^n - n(n-1)I_{n-2}, \quad n \geq 2.$$

(b) Hence find the value of  $I_4$ , giving your answer in terms of  $\pi$ .

[6]

Total: 11





8. The rectangular hyperbola  $C$  has equation  $xy = c^2$ , where  $c$  is a positive constant.

- (a) Show that an equation of the tangent to  $C$  at the point  $P\left(cp, \frac{c}{p}\right)$  is [4]

$$x + yp^2 = 2cp.$$

The tangent to  $C$  at  $P$  meets the  $x$ -axis at the point  $X$ .

The point  $Q$  on  $C$  has coordinates  $\left(cq, \frac{c}{q}\right)$ ,  $q \neq p$  such that  $QX$  is parallel to the  $y$ -axis

- (b) Show that  $q = 2p$ . [3]

$M$  is the mid-point of  $PQ$ .

- (c) Find, in Cartesian form, an equation of the locus of  $M$  as  $p$  varies. [5]

Total: 12

