## Solomon Practice Paper

## Mechanics 2B

Time allowed: 90 mintues

Centre:

Name:

Teacher:

Question	Points	Score
1	7	
2	7	
3	10	
4	10	
5	10	
6	15	
7	16	
Total:	75	

## How I can achieve better:

- •
- •
- •
- •




1.	A bullet of mass 25g is fired directly at a fixed wooden block of thickness 4cm and passes through it. When the bullet hits the block, it is travelling horizontally at 200 ms <sup>-1</sup> . The block exerts a constant resistive force of 8000N on the bullet.	
	(a) Find the work done by the block on the bullet.	[2]
	By using the Work-Energy principle,	
	(b) show that the bullet emerges from the block with speed $120 \text{ ms}^{-1}$ .	[5]
		Total: 7
		-
		-
		-
		-
		-
		-
		-
		_
		-
		-
		-
		-
		-
		-
		-
		-
		-
		-
		-



2.	A car is travelling along a straight horizontal road against resistances to motion which are constant and total 2000 N. When the engine of the car is working at a rate of $H$ kilowatts, the maximum speed of the car is $30~{\rm ms}^{-1}$ .	
	(a) Find the value of $H$ .	[3]
	The car driver wishes to overtake another vehicle so she increases the rate of working of the engine by $20\%$ and this results in an initial acceleration of $0.32~\mathrm{ms}^{-2}$ .	
	Assuming that the resistances to motion remain constant,	
	(b) find the mass of the car.	[4]
		Total: 7



3. Figure shows a uniform triangular lamina ABC placed with edge BC along the line of greatest slope of a plane inclined at an angle  $\theta$  to the horizontal.



The lengths AC and BC are 15 cm and 9 cm respectively and  $\angle ABC$  is a right angle.

(a) Find the distance of the centre of mass of the lamina from

[6]

Total: 10

- i. AB,
- ii. BC.


Assuming that the plane is rough enough to prevent the lamina from slipping,

(b) find in degrees, correct to 1 decimal place, the maximum value of  $\theta$  for which the lamina remains in equilibrium. [4]



4.	The velocity $v \text{ ms}^{-1}$ of a particle $P$ at time $t$ seconds is given by $v = 3t\mathbf{i} - t^2\mathbf{j}$ .	
	(a) Find the magnitude of the acceleration of $P$ when $t=2$ .	[4]
	When $t = 0$ , the displacement of $P$ from a fixed origin $O$ is $(6\mathbf{i} + 12\mathbf{j}) \text{ ms}^{-1}$ , where $\mathbf{i}$ and $\mathbf{j}$ are perpendicular horizontal unit vectors.	;
	(b) Show that the displacement of $P$ from $O$ when $t = 6$ is given by $k(\mathbf{i} - \mathbf{j})m$ , where $k$ is an integer which you should find.	[6]
	r	Total: 10
		-
		-
		-
		-
		-
		-
		-
		-
		-
		-
		=
		-
		-
		-
		-
		-
		-

5. A uniform rod AB of length 2a and mass 8kg is smoothly hinged to a vertical wall at A.



The rod is held in equilibrium inclined at an angle of  $20^{\circ}$  to the horizontal by a force of magnitude F newtons acting horizontally at B which is below the level of A as shown in Figure.

(a) Find, correct to 3 significant figures, the value of F.

[4] [6]

(b) Show that the magnitude of the reaction at the hinge is 133 N, correct to 3 significant figures, and find to the nearest degree the acute angle which the reaction makes with the vertical.

Total: 10



6.	A particle $P$ is projected from a point $A$ on horizontal ground with speed $u$ at an angle of elevation $\alpha$ and moves freely under gravity.		
	P hits the ground at the point $B$ .		
	(a) Show that $AB = gu2\sin(2\alpha)$ .		[6]
	An archer fires an arrow with an initial speed of 45 ms <sup>-1</sup> at a target which is level with the point of projection and at a distance of 80 m.  Given that the arrow hits the target,		
	(b) find in degrees, correct to 1 decimal place, the two possible angles of projection.		[٢]
			[5]
	(c) Write down, with a reason, which of the two possible angles of projection would give the shortest time of flight.		[2]
	(d) Show that the minimum time of flight is 1.8 seconds, correct to 1 decimal place.		[2]
	${ m T}$	otal:	15
	<u> </u>		



[7]

[9]

16

7.	A smooth sphere $A$ of mass $4m$ is moving on a smooth horizontal plane with speed $u$ . It collides directly with a stationary smooth sphere $B$ of mass $5m$ and with the same radius as $A$ .	S
	The coefficient of restitution between A and B is $\frac{1}{2}$ .	
	(a) Show that after the collision the speed of $B$ is 4 times greater than the speed of $A$ .	
	Sphere $B$ subsequently hits a smooth vertical wall at right angles. After rebounding from the wall, $B$ collides with $A$ again and as a result of this collision, $B$ comes to rest.	е
	Given that the coefficient of restitution between $B$ and the wall is e,	
	(b) find e.	Total
		Totar
		_
		_
		_
		_
		_
		_
		=
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_

