

# Solomon Practice Paper

## Mechanics 3A

**Time allowed:** 90 minutes

**Centre:** [www.CasperYC.club](http://www.CasperYC.club)


**Name:**

**Teacher:**

Question	Points	Score
1	7	
2	7	
3	10	
4	11	
5	13	
6	13	
7	14	
Total:	75	

**How I can achieve better:**

- 
- 
- 




Last updated:

July 14, 2025




1. A particle of mass  $0.6\text{kg}$  is attached to one end of a light elastic spring of natural length  $1\text{m}$  and modulus of elasticity  $30\text{N}$ . The other end of the spring is fixed to a point  $O$  which lies on a smooth plane inclined at an angle  $\alpha$  to the horizontal where  $\tan \alpha = \frac{3}{4}$  as shown in Figure.



The particle is held at rest on the slope at a point  $1.2\text{m}$  from  $O$  down the line of greatest slope of the plane.

- (a) Find the tension in the spring. [2]
- (b) Find the initial acceleration of the particle. [5]

Total: 7



2. A particle  $P$  of mass  $0.5\text{kg}$  moves along the positive  $x$ -axis under the action of a single force directed away from the origin  $O$ . When  $P$  is  $x$  metres from  $O$ , the magnitude of the force is  $3x^{\frac{1}{2}}\text{N}$  and  $P$  has a speed of  $v\text{ ms}^{-1}$ .

Given that when  $x = 1$ ,  $P$  is moving away from  $O$  with speed  $2\text{ ms}^{-1}$ ,

- (a) find an expression for  $v^2$  in terms of  $x$ , [5]
- (b) show that when  $x = 4$ ,  $P$  has a speed of  $7.7\text{ms}^{-1}$ , correct to 1 decimal place. [2]

Total: 7



3. A particle is performing simple harmonic motion along a straight line between the points  $A$  and  $B$  where  $AB = 8$  m. The period of the motion is 12 seconds.

(a) Find the maximum speed of the particle in terms of  $\pi$ . [4]

The points  $P$  and  $Q$  are on the line  $AB$  at distances of 3m and 6m respectively from  $A$ .

(b) Find, correct to 3 significant figures, the time it takes for the particle to travel directly from  $P$  to  $Q$ . [6]

Total: 10



4. Whilst in free-fall a parachutist falls vertically such that his velocity,  $v \text{ ms}^{-1}$ , when he is  $x$  metres below his initial position is given by

$$v^2 = kg \left( 1 - e^{-\frac{2x}{k}} \right),$$

where  $k$  is a constant.

Given that he experiences an acceleration of  $f \text{ ms}^{-2}$ ,

- (a) show that  $f = ge^{-\frac{2x}{k}}$ . [4]

After falling a large distance, his velocity is constant at  $49 \text{ ms}^{-1}$ .


- (b) Find the value of  $k$ . [3]

- (c) Hence, express  $f$  in the form  $(\lambda - \mu v^2)$  where  $\lambda$  and  $\mu$  are constants which you should find. [4]

Total: 11



5. A firework is modelled as a uniform solid formed by joining the plane surface of a right circular cone of height  $2r$  and base radius  $r$ , to one of the plane surfaces of a cylinder of height  $h$  and base radius  $r$  as shown in Figure.



Using this model,

- (a) show that the distance of the centre of mass of the firework from its plane base is [9]


The firework is to be launched from rough ground inclined at an angle  $\alpha$  to the horizontal. Given that the firework does not slip or topple and that  $h = 4r$ ,

- (b) Find, correct to the nearest degree, the maximum value of  $\alpha$ . [4]

Total: 13



6. The two ends of a light inextensible string of length  $3a$  are attached to fixed points  $Q$  and  $R$  which are a distance of  $a\sqrt{3}$  apart with  $R$  vertically below  $Q$ . A particle  $P$  of mass  $m$  is attached to the string at a distance of  $2a$  from  $Q$ .



$P$  is given a horizontal speed,  $u$ , such that it moves in a horizontal circle with both sections of the string taut as shown in Figure.

- (a) Show that  $\angle PRQ$  is a right angle. [2]
- (b) Find  $\angle PQR$  in degrees. [1]
- (c) Find, in terms of  $a, g, m$  and  $u$ , the tension in the section of string [7]
  - i.  $PQ$ ,
  - ii.  $PR$ .
- (d) Show that  $u^2 \geq \frac{ga}{\sqrt{3}}$ . [3]

Total: 13



7. A particle of mass 2kg is attached to one end of a light elastic string of natural length 1m and modulus of elasticity 50N. The other end of the string is attached to a fixed point  $O$  on a rough horizontal plane and the coefficient of friction between the particle and the plane is  $\frac{10}{49}$ .

The particle is projected from  $O$  along the plane with an initial speed of  $5 \text{ ms}^{-1}$ .

(a) Show that the greatest distance from  $O$  which the particle reaches is 1.84 m. [9]

(b) Find, correct to 2 significant figures, the speed at which the particle returns to  $O$ . [5]

Total: 14

