

# Solomon Practice Paper

## Pure Mathematics 5C

Time allowed: 90 minutes

Centre: [www.CasperYC.club](http://www.CasperYC.club)

Name:

Teacher:


Question	Points	Score
1	5	
2	7	
3	10	
4	12	
5	12	
6	13	
7	16	
Total:	75	

How I can achieve better:

- 
- 
- 



Last updated: May 5, 2023



1. The curve  $C$  has intrinsic equation [5]

$$s = 4 \sec^3(\psi), \quad 0 \leq \psi < \frac{\pi}{2}.$$

Find the radius of curvature of  $C$  at the point where  $\psi = \frac{\pi}{4}$ .

2. Solve the equation [7]

$$5 \coth(x) + 1 = 7 \operatorname{cosech}(x),$$

giving your answer in terms of natural logarithms.

3. (a) Show that [3]

$$\frac{d}{dx} \arccos(x) = -\frac{1}{\sqrt{1-x^2}}.$$

- (b) The curve with equation [7]

$$y = \arccos(x) - \frac{1}{2} \ln(1-x^2), \quad -1 < x < 1,$$

has a stationary point in the interval  $0 < x < 1$ .

Find the exact coordinates of this stationary point.

Total: 10

4. (a) Express  $3 - 6x - 9x^2$  in the form  $a - (bx + c)^2$  where  $a, b$  and  $c$  are constants. [2]

Hence, or otherwise, find

- (b) [4]

$$\int \frac{1}{\sqrt{3-6x-9x^2}} dx,$$

- (c) expressing your answer to part (b) in terms of natural logarithms. [6]

$$\int_{-\frac{1}{3}}^0 \frac{1}{\sqrt{3-6x-9x^2}} dx,$$

Total: 12


- 5.

$$f(x) = \operatorname{arctanh}\left(\frac{x^2-1}{x^2+1}\right), \quad x > 0.$$

- (a) Using the definitions of  $\sinh(x)$  and  $\cosh(x)$  in terms of exponentials, express  $\tanh(x)$  in terms of  $e^x$  and  $e^{-x}$ . [1]

- (b) Hence prove that  $f(x) = \ln(x)$ . [6]

- (c) Hence, or otherwise, show that the area bounded by the curve  $y = \operatorname{arctanh}\left(\frac{x^2-1}{x^2+1}\right)$ , the positive  $x$ -axis and the line  $x = 2e$  is  $2e \ln(2) + 1$ . [5]



6. The ellipse  $C$  has equation

$$\frac{x^2}{25} + \frac{y^2}{9} = 1.$$

(a) Find an equation of the normal to  $C$  at the point  $P(5 \cos(\theta), 3 \sin(\theta))$ . [5]

The normal to  $C$  at  $P$  meets the coordinate axes at  $Q$  and  $R$ .

Given that  $ORSQ$  is a rectangle, where  $O$  is the origin,

(b) show that, as  $\theta$  varies, the locus of  $S$  is an ellipse and find its equation in Cartesian form. [8]

Total: 13

7.

$$I_n(x) = \int_0^x \cos^n(2t) dt, \quad n \geq 0.$$

(a) Show that [7]


$$nI_n(x) = \frac{1}{2} \sin(2x) \cos^{n-1}(2x) + (n-1)I_{n-2}(x), \quad n \neq 2.$$

(b) Find  $I_0\left(\frac{\pi}{4}\right)$  in terms of  $\pi$ . [2]

Figure shows the curve with polar equation

$$r = a \cos^2(2\theta), \quad 0 \leq \theta \leq \frac{\pi}{4},$$

where  $a$  is a positive constant.



(c) Using your answers to parts (a) and (b), or otherwise, calculate the area bounded by the curve and the half-lines  $\theta = 0$  and  $\theta = \frac{\pi}{4}$ . [7]

Total: 16

