

# Solomon Practice Paper

## Statistics S1 – G

**Time allowed:** 90 minutes

**Centre:** [www.CasperYC.club](http://www.CasperYC.club)


**Name:**

**Teacher:**

Question	Points	Score
1	7	
2	8	
3	9	
4	11	
5	11	
6	14	
7	15	
Total:	75	

**How I can achieve better:**

- 
- 
- 



Last updated:

July 14, 2025




1. The discrete random variable  $Y$  has the following probability distribution.

$y$	-2	-1	0	1	2
$\Pr(Y = y)$	0.1	0.15	0.2	0.3	0.25

Find

- (a)  $F(0.5)$ , [1]
- (b)  $P(-1 < Y < 1.9)$ , [2]
- (c)  $E(Y)$ , [2]
- (d)  $E(3Y - 1)$ . [2]

Total: 7



2. A supermarket manager believes that those of her staff on lower rates of pay tend to work more hours of overtime.

- (a) A supermarket manager believes that those of her staff on lower rates of pay tend to work more hours of overtime. [1]

To investigate her theory the manager recorded the number of hours of overtime,  $h$ , worked by each of the store's 18 full-time staff during one week. She also recorded each employee's hourly rate of pay,  $\mathcal{L}p$ , and summarised her results as follows:

$$\sum p = 86, \quad \sum h = 104.5, \quad \sum p^2 = 420.58, \quad \sum h^2 = 830.25, \quad \sum ph = 487.3$$

- (b) Calculate the product moment correlation coefficient for these data. [5]

- (c) Comment on the manager's hypothesis. [2]

Total: 8




The data is grouped and coded using  $y = \frac{C - 3250}{250}$ , where  $C$  is the mid-point in pounds of each class, giving

(a) Using these values, calculate estimates of the mean and standard deviation of the cost of the receptions in the sample. [6]

The median of the data was £3050.

(c) Comment on the skewness of the data and suggest a reason for it. [2]

Total: 9



4. The random variable  $A$  is normally distributed with a mean of 32.5 and a variance of 18.6 Find

(a)  $P(A < 38.2)$ , [3]


(b)  $P(31 \leq A \leq 35)$ , [4]

The random variable  $B$  is normally distributed with a standard deviation of 7.2.

Given also that  $P(B > 110) = 0.138$ ,

(c) find the mean of  $B$ . [4]

Total: 11



- |                    |    |    |    |   |
|--------------------|----|----|----|---|
| Number of attempts | 1  | 2  | 3  | 4 |
| Number of children | 43 | 26 | 13 | 3 |

- $$P(X = x) = \begin{cases} k(20 - x^2), & x = 1, 2, 3, 4 \\ 0, & \text{otherwise} \end{cases}$$

- Total: 11



6. Serving against his regular opponent, a tennis player has a 65% chance of getting his first serve in. If his first serve is in he then has a 70% chance of winning the point but if his first serve is not in, he only has a 45% chance of winning the point.

(a) Represent this information on a tree diagram.

[3]

For a point on which this player served to his regular opponent, find the probability that

(b) he won the point,

[3]

(c) his first serve went in given that he won the point,

[3]

(d) his first serve didn't go in given that he lost the point.

[5]

Total: 14



$l$ (metres)	0.5	0.8	1.0	1.5	2	4	6
$P$ (£)	2.50	3.40	4.00	5.20	6.00	10.50	15.00

[3]

$$\sum l = 15.8, \quad \sum P = 46.6, \quad \sum l^2 = 60.14, \quad \sum lP = 159.77$$

[7]

$$[1]$$

[2]

[2]

