Solomon Practice Paper

Statistics S2 - A

Time allowed:	90	minutes
---------------	----	---------

Name:

Teacher:

Question	Points	Score
1	5	
2	8	
3	10	
4	10	
5	11	
6	12	
7	19	
Total:	75	

How I can achieve better:

- •



Last updated: July 14, 2025



١

a be [
ſ
[
Total:
00
\sim

2. The continuous random variable X has the following cumulative distribution function:

$$\mathbf{F}(x) = \begin{cases} 0, & x < 0, \\ \frac{1}{64}(16x - x^2), & 0 \le x \le 8, \\ 1, & x > 8. \end{cases}$$

- (a) Find $\Pr(X > 5)$.
- (b) Find and specify fully the probability density function f(x) of X.
- (c) Sketch f(x) for all values of x.

Total: 8

[2]

[3]

[3]



- 3. An electrician records the number of repairs of different types of appliances that he makes each day. His records show that over 40 working days he repaired a total of 180 CD players.
 - (a) Explain why a Poisson distribution may be suitable for modelling the number of CD players [4] he repairs each day and find the parameter for this distribution.
 - (b) Find the probability that on one particular day he repairs
 - i. no CD players,
 - ii. more than 6 CD players.
 - (c) Find the probability that over 10 working days he will repair more than 6 CD players on [3] exactly 3 of the days.

Total: 10

[3]



- 4. A teacher wants to investigate the sports played by students at her school in their free time. She decides to ask a random sample of 120 pupils to complete a short questionnaire.
 - (a) Give two reasons why the teacher might choose to use a sample survey rather than a census. [2]
 - (b) Suggest a suitable sampling frame that she could use.

The teacher believes that 1 in 20 of the students play tennis in their free time. She uses the data collected from her sample to test if the proportion is different from this.

- (c) Using a suitable approximation and stating the hypotheses that she should use, find the [6] critical region for this test. The probability for each tail of the region should be as close as possible to 5%.
- (d) State the significance level of this test.

[1]

[1]

Total: 10



5. As part of a business studies project, 8 groups of students are each randomly allocated 10 different shares from a listing of over 300 share prices in a newspaper. Each group has to follow the changes in the price of their shares over a 3–month period.

At the end of the 3 months, 35% of all the shares in the listing have increased in price and the rest have decreased.

- (a) Find the probability that, for the 10 shares of one group,
 - i. exactly 6 have gone up in price,
 - ii. more than 5 have gone down in price.
- (b) Using a suitable approximation, find the probability that of the 80 shares allocated in total
 to the groups, more than 55 will have decreased in value.

Total: 11

[5]



- 6. A shoe shop sells on average 4 pairs of shoes per hour on a weekday morning.
 - (a) Suggest a suitable distribution for modelling the number of sales made per hour on a weekday [1] morning and state the value of any parameters needed.
 - (b) Explain why this model might have to be modified for modelling the number of sales made [1] per hour on a Saturday morning.
 - (c) Find the probability that on a weekday morning the shop sells
 - i. more than 4 pairs in a one–hour period,

ii. no pairs in a half–hour period,

iii. more than 4 pairs during each hour from 9 am until noon.

The area manager visits the shop on a weekday morning, the day after an advert appears in a local paper. In a one-hour period the shop sells 7 pairs of shoes, leading the manager to believe that the advert has increased the shop's sales.

(d) Stating your hypotheses clearly, test at the 5% level of significance whether or not there is [4] evidence of an increase in sales following the appearance of the advert.

Total: 12

[6]



7. The continuous random variable T has the following probability density function:

$$\mathbf{f}(t) = \begin{cases} k(t^2 + 2), & 0 \le t \le 3, \\ 0, & \text{otherwise.} \end{cases}$$

(a)	Show that	[4]
(b)	Sketch $f(t)$ for all values of t .	[3]
(c)	State the mode of T .	[1]
(d)	Find $E(T)$.	[5]
(e)	Show that the standard deviation of T is 0.798 correct to 3 significant figures.	[6]

Total: 19

www.CasperYC.club