


THERE ARE 13 QUESTIONS IN THIS PAPER.

Q1 Q2 Q3 Q4 Q5 Q6 Q7 Q8 Q9 Q10 Q11 Q12 Q13

Sixth Term Examination Paper

11-S2



Compiled by: Dr Yu 郁博士

www.CasperYC.club

Last updated: May 8, 2025



SUGGESTIONS TO DRYUFROMSHANGHAI@QQ.COM

Section A: Pure Mathematics

- 1 (i) Sketch the curve $y = \sqrt{1-x} + \sqrt{3+x}$. Use your sketch to show that only one real value of x satisfies

$$\sqrt{1-x} + \sqrt{3+x} = x + 1,$$

and give this value.

- (ii) Determine graphically the number of real values of x that satisfy

$$2\sqrt{1-x} = \sqrt{3+x} + \sqrt{3-x}.$$

Solve this equation.

- 2 Write down the cubes of the integers 1, 2, ..., 10. The positive integers x, y and z , where $x < y$, satisfy

$$x^3 + y^3 = kz^3, \quad (*)$$

where k is a given positive integer.

- (i) In the case $x + y = k$, show that

$$z^3 = k^2 - 3kx + 3x^2.$$

Deduce that $(4z^3 - k^2)/3$ is a perfect square and that $\frac{1}{4}k^2 \leq z^3 < k^2$. Use these results to find a solution of $(*)$ when $k = 20$.

- (ii) By considering the case $x + y = z^2$, find two solutions of $(*)$ when $k = 19$.

3 In this question, you may assume without proof that any function f for which $f'(x) \geq 0$ is *increasing*; that is, $f(x_2) \geq f(x_1)$ if $x_2 \geq x_1$.

(i) (a) Let $f(x) = \sin x - x \cos x$. Show that $f(x)$ is increasing for $0 \leq x \leq \frac{1}{2}\pi$ and deduce that $f(x) \geq 0$ for $0 \leq x \leq \frac{1}{2}\pi$.

(b) Given that $\frac{d}{dx}(\arcsin x) \geq 1$ for $0 \leq x < 1$, show that

$$\arcsin x \geq x \quad (0 \leq x < 1).$$

(c) Let $g(x) = x \operatorname{cosec} x$ for $0 < x < \frac{1}{2}\pi$. Show that g is increasing and deduce that

$$(\arcsin x) x^{-1} \geq x \operatorname{cosec} x \quad (0 < x < 1).$$

(ii) Given that $\frac{d}{dx}(\arctan x) \leq 1$ for $x \geq 0$, show by considering the function $x^{-1} \tan x$ that

$$(\tan x)(\arctan x) \geq x^2 \quad (0 < x < \frac{1}{2}\pi).$$

4 (i) Find all the values of θ , in the range $0^\circ < \theta < 180^\circ$, for which $\cos \theta = \sin 4\theta$. Hence show that

$$\sin 18^\circ = \frac{1}{4}(\sqrt{5} - 1).$$

(ii) Given that

$$4 \sin^2 x + 1 = 4 \sin^2 2x,$$

find all possible values of $\sin x$, giving your answers in the form $p + q\sqrt{5}$ where p and q are rational numbers.

(iii) Hence find two values of α with $0^\circ < \alpha < 90^\circ$ for which

$$\sin^2 3\alpha + \sin^2 5\alpha = \sin^2 6\alpha.$$

- 5** The points A and B have position vectors \mathbf{a} and \mathbf{b} with respect to an origin O , and O, A and B are non-collinear. The point C , with position vector \mathbf{c} , is the reflection of B in the line through O and A .

- (i) Show that \mathbf{c} can be written in the form

$$\mathbf{c} = \lambda \mathbf{a} - \mathbf{b}$$

$$\text{where } \lambda = \frac{2 \mathbf{a} \cdot \mathbf{b}}{\mathbf{a} \cdot \mathbf{a}}.$$

- (ii) The point D , with position vector \mathbf{d} , is the reflection of C in the line through O and B . Show that \mathbf{d} can be written in the form

$$\mathbf{d} = \mu \mathbf{b} - \lambda \mathbf{a}$$

for some scalar μ to be determined.

- (iii) Given that A, B and D are collinear, find the relationship between λ and μ . In the case $\lambda = -\frac{1}{2}$, determine the cosine of $\angle AOB$ and describe the relative positions of A, B and D .

- 6** For any given function f , let

$$I = \int [f'(x)]^2 [f(x)]^n dx, \quad (*)$$

where n is a positive integer. Show that, if $f(x)$ satisfies $f''(x) = kf(x)f'(x)$ for some constant k , then $(*)$ can be integrated to obtain an expression for I in terms of $f(x), f'(x), k$ and n .

- (i) Verify your result in the case $f(x) = \tan x$. Hence find

$$\int \frac{\sin^4 x}{\cos^8 x} dx.$$

- (ii) Find

$$\int \sec^2 x (\sec x + \tan x)^6 dx.$$

- 7 The two sequences a_0, a_1, a_2, \dots and b_0, b_1, b_2, \dots have general terms

$$a_n = \lambda^n + \mu^n \quad \text{and} \quad b_n = \lambda^n - \mu^n,$$

respectively, where $\lambda = 1 + \sqrt{2}$ and $\mu = 1 - \sqrt{2}$.

- (i) Show that $\sum_{r=0}^n b_r = -\sqrt{2} + \frac{1}{\sqrt{2}} a_{n+1}$, and give a corresponding result for $\sum_{r=0}^n a_r$.

- (ii) Show that, if n is odd,

$$\sum_{m=0}^{2n} \left(\sum_{r=0}^m a_r \right) = \frac{1}{2} b_{n+1}^2,$$

and give a corresponding result when n is even.

- (iii) Show that, if n is even,

$$\left(\sum_{r=0}^n a_r \right)^2 - \sum_{r=0}^n a_{2r+1} = 2,$$


and give a corresponding result when n is odd.

- 8** The end A of an inextensible string AB of length π is attached to a point on the circumference of a fixed circle of unit radius and centre O . Initially the string is straight and tangent to the circle. The string is then wrapped round the circle until the end B comes into contact with the circle. The string remains taut during the motion, so that a section of the string is in contact with the circumference and the remaining section is straight.

- (i) Taking O to be the origin of cartesian coordinates with A at $(-1, 0)$ and B initially at $(-1, \pi)$, show that the curve described by B is given parametrically by

$$x = \cos t + t \sin t \quad \text{and} \quad y = \sin t - t \cos t,$$

where t is the angle shown in the diagram.



Find the value, t_0 , of t for which x takes its maximum value on the curve, and sketch the curve.

- (ii) Use the area integral $\int y \frac{dx}{dt} dt$ to find the area between the curve and the x axis for $\pi \geq t \geq t_0$.
- (iii) Find the area swept out by the string (that is, the area between the curve described by B and the semicircle shown in the diagram).

Section B: Mechanics

- 9 Two particles, A of mass $2m$ and B of mass m , are moving towards each other in a straight line on a smooth horizontal plane, with speeds $2u$ and u respectively. They collide directly.

- (i) Given that the coefficient of restitution between the particles is e , where $0 < e \leq 1$, determine the speeds of the particles after the collision.
- (ii) After the collision, B collides directly with a smooth vertical wall, rebounding and then colliding directly with A for a second time. The coefficient of restitution between B and the wall is f , where $0 < f \leq 1$. Show that the velocity of B after its second collision with A is

$$\frac{2}{3}(1 - e^2)u - \frac{1}{3}(1 - 4e^2)fu$$


towards the wall and that B moves towards (not away from) the wall for all values of e and f .

- 10 (i) A particle is projected from a point on a horizontal plane, at speed u and at an angle θ above the horizontal. Let H be the maximum height of the particle above the plane. Derive an expression for H in terms of u , g and θ .
- (ii) A particle P is projected from a point O on a smooth horizontal plane, at speed u and at an angle θ above the horizontal. At the same instant, a second particle R is projected horizontally from O in such a way that R is vertically below P in the ensuing motion. A light inextensible string of length $\frac{1}{2}H$ connects P and R . Show that the time that elapses before the string becomes taut is

$$(\sqrt{2} - 1)\sqrt{H/g}.$$

- (iii) When the string becomes taut, R leaves the plane, the string remaining taut.
- (iv) Given that P and R have equal masses, determine the total horizontal distance, D , travelled by R from the moment its motion begins to the moment it lands on the plane again, giving your answer in terms of u , g and θ . Given that $D = H$, find the value of $\tan \theta$.

- 11** Three non-collinear points A, B and C lie in a horizontal ceiling. A particle P of weight W is suspended from this ceiling by means of three light inextensible strings AP, BP and CP , as shown in the diagram. The point O lies vertically above P in the ceiling.



The angles AOB and AOC are $90^\circ + \theta$ and $90^\circ + \phi$, respectively, where θ and ϕ are acute angles such that $\tan \theta = \sqrt{2}$ and $\tan \phi = \frac{1}{4}\sqrt{2}$. The strings AP, BP and CP make angles $30^\circ, 90^\circ - \theta$ and 60° , respectively, with the vertical, and the tensions in these strings have magnitudes T, U and V respectively.

- (i)** Show that the unit vector in the direction PB can be written in the form

$$-\frac{1}{3}\mathbf{i} - \frac{\sqrt{2}}{3}\mathbf{j} + \frac{\sqrt{2}}{\sqrt{3}}\mathbf{k},$$

where \mathbf{i}, \mathbf{j} and \mathbf{k} are the usual mutually perpendicular unit vectors with \mathbf{j} parallel to OA and \mathbf{k} vertically upwards.

- (ii)** Find expressions in vector form for the forces acting on P .

- (iii)** Show that $U = \sqrt{6}V$ and find T, U and V in terms of W .

Section C: Probability and Statistics

- 12** Xavier and Younis are playing a match. The match consists of a series of games and each game consists of three points. Xavier has probability p and Younis has probability $1 - p$ of winning the first point of any game. In the second and third points of each game, the player who won the previous point has probability p and the player who lost the previous point has probability $1 - p$ of winning the point. If a player wins two consecutive points in a single game, the match ends and that player has won; otherwise the match continues with another game.

- (i) Let w be the probability that Younis wins the match. Show that, for $p \neq 0$,

$$w = \frac{1 - p^2}{2 - p}.$$

Show that $w > \frac{1}{2}$ if $p < \frac{1}{2}$, and $w < \frac{1}{2}$ if $p > \frac{1}{2}$. Does w increase whenever p decreases?

- (ii) If Xavier wins the match, Younis gives him £1;
if Younis wins the match, Xavier gives him £ k .
Find the value of k for which the game is 'fair' in the case when $p = \frac{2}{3}$.
- (iii) What happens when $p = 0$?

13 What property of a distribution is measured by its *skewness*?

(i) One measure of skewness, γ , is given by

$$\gamma = \frac{E((X - \mu)^3)}{\sigma^3},$$

where μ and σ^2 are the mean and variance of the random variable X . Show that

$$\gamma = \frac{E(X^3) - 3\mu\sigma^2 - \mu^3}{\sigma^3}.$$

The continuous random variable X has probability density function f where

$$f(x) = \begin{cases} 2x & \text{for } 0 \leq x \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Show that for this distribution $\gamma = -\frac{2\sqrt{2}}{5}$.

(ii) The *decile skewness*, D , of a distribution is defined by

$$D = \frac{F^{-1}(\frac{9}{10}) - 2F^{-1}(\frac{1}{2}) + F^{-1}(\frac{1}{10})}{F^{-1}(\frac{9}{10}) - F^{-1}(\frac{1}{10})},$$

where F^{-1} is the inverse of the cumulative distribution function. Show that, for the above distribution, $D = 2 - \sqrt{5}$. The *Pearson skewness*, P , of a distribution is defined by

$$P = \frac{3(\mu - M)}{\sigma},$$

where M is the median. Find P for the above distribution and show that $D > P > \gamma$.