


THERE ARE 12 QUESTIONS IN THIS PAPER.

Q1 Q2 Q3 Q4 Q5 Q6 Q7 Q8 Q9 Q10 Q11 Q12

## Sixth Term Examination Paper


20-S2



**Compiled by: Dr Yu 郁博士**

[www.CasperYC.club](http://www.CasperYC.club)

**Last updated: May 8, 2025**



SUGGESTIONS TO [DRYUFROMSHANGHAI@QQ.COM](mailto:DRYUFROMSHANGHAI@QQ.COM)

## Section A: Pure Mathematics

- 1 (i) Use the substitution  $x = \frac{1}{1-u}$ , where  $0 < u < 1$ , to find in terms of  $x$  the integral

$$\int \frac{1}{x^{\frac{3}{2}}(x-1)^{\frac{1}{2}}} dx \quad (\text{where } x > 1).$$

- (ii) Find in terms of  $x$  the integral

$$\int \frac{1}{(x-2)^{\frac{3}{2}}(x+1)^{\frac{1}{2}}} dx \quad (\text{where } x > 2).$$

- (iii) Show that

$$\int_2^{\infty} \frac{1}{(x-1)(x-2)^{\frac{1}{2}}(3x-2)^{\frac{1}{2}}} dx = \frac{1}{3}\pi.$$

- 2 The curves  $C_1$  and  $C_2$  both satisfy the differential equation

$$\frac{dy}{dx} = \frac{kxy - y}{x - kxy},$$

where  $k = \ln 2$ .

All points on  $C_1$  have positive  $x$  and  $y$  co-ordinates and  $C_1$  passes through  $(1, 1)$ . All points on  $C_2$  have negative  $x$  and  $y$  co-ordinates and  $C_2$  passes through  $(-1, -1)$ .

- (i) Show that the equation of  $C_1$  can be written as

$$(x-y)^2 = (x+y)^2 - 2^{x+y}.$$

Determine a similar result for curve  $C_2$ .

Hence show that  $y = x$  is a line of symmetry of each curve.

- (ii) Sketch on the same axes the curves  $y = x^2$  and  $y = 2^x$ , for  $x \geq 0$ .  
Hence show that  $C_1$  lies between the lines  $x + y = 2$  and  $x + y = 4$ .

Sketch curve  $C_1$ .

- (iii) Sketch curve  $C_2$ .

- 3 A sequence  $u_1, u_2, \dots, u_n$  of positive real numbers is said to be unimodal if there is a value  $k$  such that

$$u_1 \leq u_2 \leq \dots \leq u_k$$

and

$$u_k \geq u_{k+1} \geq \dots \geq u_n.$$

So the sequences 1, 2, 3, 2, 1; 1, 2, 3, 4, 5; 1, 1, 3, 3, 2 and 2, 2, 2, 2, 2 are all unimodal, but 1, 2, 1, 3, 1 is not.

A sequence  $u_1, u_2, \dots, u_n$  of positive real numbers is said to have property  $L$  if

$$u_{r-1}u_{r+1} \leq u_r^2$$

for all  $r$  with  $2 \leq r \leq n-1$ .

- (i) Show that, in any sequence of positive real numbers with property  $L$ ,

$$u_{r-1} \geq u_r \implies u_r \geq u_{r+1}.$$

Prove that any sequence of positive real numbers with property  $L$  is unimodal.

- (ii) A sequence  $u_1, u_2, \dots, u_n$  of real numbers satisfies

$$u_r = 2\alpha u_{r-1} - \alpha^2 u_{r-2}$$

for  $3 \leq r \leq n$ , where  $\alpha$  is a positive real constant.

Prove that, for  $2 \leq r \leq n$ ,

$$u_r - \alpha u_{r-1} = \alpha^{r-2} (u_2 - \alpha u_1)$$

and, for  $2 \leq r \leq n-1$ ,

$$u_r^2 - u_{r-1}u_{r+1} = (u_r - \alpha u_{r-1})^2.$$

Hence show that the sequence consists of positive terms and is unimodal, provided  $u_2 > \alpha u_1 > 0$ .

In the case  $u_1 = 1$  and  $u_2 = 2$ , prove by induction that

$$u_r = (2-r)\alpha^{r-1} + 2(r-1)\alpha^{r-2}.$$

Let  $\alpha = 1 - \frac{1}{N}$ , where  $N$  is an integer with  $2 \leq N \leq n$ .

In the case  $u_1 = 1$  and  $u_2 = 2$ , prove that  $u_r$  is largest when  $r = N$ .

4 (i) Given that  $a, b$  and  $c$  are the lengths of the sides of a triangle, explain why  $c < a + b, a < b + c$  and  $b < a + c$ .

(ii) Use a diagram to show that the converse of the result in part (i) also holds: if  $a, b$  and  $c$  are positive numbers such that  $c < a + b, a < b + c$  and  $b < c + a$  then it is possible to construct a triangle with sides of length  $a, b$  and  $c$ .

(iii) When  $a, b$  and  $c$  are the lengths of the sides of a triangle, determine in each case whether the following sets of three lengths can

- always
- sometimes but not always
- never

form the sides of a triangle. Prove your claims.

(a)  $a + 1, b + 1, c + 1$ .

(b)  $\frac{a}{b}, \frac{b}{c}, \frac{c}{a}$ .

(c)  $|a - b|, |b - c|, |c - a|$ .

(d)  $a^2 + bc, b^2 + ca, c^2 + ab$ .

(iv) Let  $f$  be a function defined on the positive real numbers and such that, whenever  $x > y > 0$ ,

$$f(x) > f(y) > 0 \quad \text{but} \quad \frac{f(x)}{x} < \frac{f(y)}{y}.$$

Show that, whenever  $a, b$  and  $c$  are the lengths of the sides of a triangle, then  $f(a), f(b)$  and  $f(c)$  can also be the lengths of the sides of a triangle.

- 5** If  $x$  is a positive integer, the value of the function  $d(x)$  is the sum of the digits of  $x$  in base 10. For example,  $d(249) = 2 + 4 + 9 = 15$ .

An  $n$ -digit positive integer  $x$  is written in the form

$$\sum_{r=0}^{n-1} a_r \times 10^r,$$

where  $0 \leq a_r \leq 9$  for all  $0 \leq r \leq n-1$  and  $a_{n-1} > 0$ .

**(i)** Prove that  $x - d(x)$  is non-negative and divisible by 9.

**(ii)** Prove that  $x - 44d(x)$  is a multiple of 9 if and only if  $x$  is a multiple of 9.

Suppose that  $x = 44d(x)$ .

Show that if  $x$  has  $n$  digits, then  $x \leq 396n$  and  $x \geq 10^{n-1}$ , and hence that  $n \leq 4$ .

Find a value of  $x$  for which  $x = 44d(x)$ .

Show that there are no further values of  $x$  satisfying this equation.

**(iii)** Find a value of  $x$  for which  $x = 107d(d(x))$ . Show that there are no further values of  $x$  satisfying this equation.

- 6** A  $2 \times 2$  matrix  $\mathbf{M}$  is real if it can be written as  $\mathbf{M} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ , where  $a, b, c$  and  $d$  are real.

In this case, the *trace* of matrix  $\mathbf{M}$  is defined to be  $\text{tr}(\mathbf{M}) = a + d$  and  $\det(\mathbf{M})$  is the determinant of matrix  $\mathbf{M}$ . In this question,  $\mathbf{M}$  is a real  $2 \times 2$  matrix.

**(i)** Prove that

$$\text{tr}(\mathbf{M}^2) = \text{tr}(\mathbf{M})^2 - 2\det(\mathbf{M}).$$

**(ii)** Prove that

$$\mathbf{M}^2 = \mathbf{I} \quad \text{but} \quad \mathbf{M} \neq -\mathbf{I} \quad \implies \quad \text{tr}(\mathbf{M}) = 0 \quad \text{and} \quad \det(\mathbf{M}) = -1,$$

and that

$$\mathbf{M}^2 = -\mathbf{I} \quad \implies \quad \text{tr}(\mathbf{M}) = 0 \quad \text{and} \quad \det(\mathbf{M}) = 1.$$

**(iii)** Use part **(ii)** to prove that

$$\mathbf{M}^4 = \mathbf{I} \quad \implies \quad \mathbf{M}^2 = \pm \mathbf{I}.$$

Find a necessary and sufficient condition on  $\det(\mathbf{M})$  and  $\text{tr}(\mathbf{M})$  so that  $\mathbf{M}^4 = -\mathbf{I}$ .

**(iv)** Give an example of a matrix  $\mathbf{M}$  for which  $\mathbf{M}^8 = \mathbf{I}$ , but which does not represent a rotation or reflection. [Note that the matrices  $\pm \mathbf{I}$  are both rotations.]

7 In this question,

$$w = \frac{2}{z - 2}.$$

- (i) Let  $z$  be the complex number  $3 + ti$ , where  $t \in \mathbb{R}$ . Show that  $|w - 1|$  is independent of  $t$ . Hence show that, if  $z$  is a complex number on the line  $\operatorname{Re}(z) = 3$  in the Argand diagram, then  $w$  lies on a circle in the Argand diagram with centre 1.

Let  $V$  be the line  $\operatorname{Re}(z) = p$ , where  $p$  is a real constant not equal to 2. Show that, if  $z$  lies on  $V$ , then  $w$  lies on a circle whose centre and radius you should give in terms of  $p$ . For which  $z$  on  $V$  is  $\operatorname{Im}(w) > 0$ ?

- (ii) Let  $H$  be the line  $\operatorname{Im}(z) = q$ , where  $q$  is a non-zero real constant. Show that, if  $z$  lies on  $H$ , then  $w$  lies on a circle whose centre and radius you should give in terms of  $q$ . For which  $z$  on  $H$  is  $\operatorname{Re}(w) > 0$ ?

8 In this question,  $f(x)$  is a quartic polynomial where the coefficient of  $x^4$  is equal to 1, and which has four real roots, 0,  $a$ ,  $b$  and  $c$ , where  $0 < a < b < c$ .

$F(x)$  is defined by  $F(x) = \int_0^x f(t) \, dt$ .

The area enclosed by the curve  $y = f(x)$  and the  $x$ -axis between 0 and  $a$  is equal to that between  $b$  and  $c$ , and half that between  $a$  and  $b$ .

- (i) Sketch the curve  $y = F(x)$ , showing the  $x$  co-ordinates of its turning points.

Explain why  $F(x)$  must have the form

$$F(x) = \frac{1}{5}x^2(x - c)^2(x - h),$$

where  $0 < h < c$ .

Find, in factorised form, an expression for  $F(x) + F(c - x)$  in terms of  $c$ ,  $h$  and  $x$ .

- (ii) If  $0 \leq x \leq c$ , explain why  $F(b) + F(x) \geq 0$  and why  $F(b) + F(x) > 0$  if  $x \neq a$ . Hence show that  $c - b = a$  or  $c > 2h$ .

By considering also  $F(a) + F(x)$ , show that  $c = a + b$  and that  $c = 2h$ .

- (iii) Find an expression for  $f(x)$  in terms of  $c$  and  $x$  only.

Show that the points of inflection on  $y = f(x)$  lie on the  $x$ -axis.

## Section B: Mechanics

- 9** Point  $A$  is a distance  $h$  above ground level and point  $N$  is directly below  $A$  at ground level. Point  $B$  is also at ground level, a distance  $d$  horizontally from  $N$ . The angle of elevation of  $A$  from  $B$  is  $\beta$ . A particle is projected horizontally from  $A$ , with initial speed  $V$ . A second particle is projected from  $B$  with speed  $U$  at an acute angle  $\theta$  above the horizontal. The horizontal components of the velocities of the two particles are in opposite directions. The two particles are projected simultaneously, in the vertical plane through  $A$ ,  $N$  and  $B$ .

Given that the two particles collide, show that

$$d \sin \theta - h \cos \theta = \frac{Vh}{U}$$

and also that

(i)  $\theta > \beta$ ;

(ii)  $U \sin \theta \geq \sqrt{\frac{gh}{2}}$ ;

(iii)  $\frac{U}{V} > \sin \beta$ .

Show that the particles collide at a height greater than  $\frac{1}{2}h$  if and only if the particle projected from  $B$  is moving upwards at the time of collision.

- 10** A particle  $P$  of mass  $m$  moves freely and without friction on a wire circle of radius  $a$ , whose axis is horizontal. The highest point of the circle is  $H$ , the lowest point of the circle is  $L$  and angle  $PHL = \theta$ . A light spring of modulus of elasticity  $\lambda$  is attached to  $P$  and to  $H$ . The natural length of the spring is  $l$ , which is less than the diameter of the circle.

- (i) Show that, if there is an equilibrium position of the particle at  $\theta = \alpha$ , where  $\alpha > 0$ , then

$$\cos \alpha = \frac{\lambda l}{2(\alpha \lambda - mgl)}.$$

Show also that there will only be such an equilibrium position if  $\lambda > \frac{2mgl}{2a - l}$ .

When the particle is at the lowest point  $L$  of the circular wire, it has speed  $u$ .

- (ii) Show that, if the particle comes to rest before reaching  $H$ , it does so when  $\theta = \beta$ , where  $\cos \beta$  satisfies

$$(\cos \alpha - \cos \beta)^2 = (1 - \cos \alpha)^2 + \frac{mu^2}{2a\lambda} \cos \alpha,$$

where  $\cos \alpha = \frac{\lambda l}{2(a\lambda - mgl)}$ .

Show also that this will only occur if

$$u^2 < \frac{2a\lambda}{m} (2 - \sec \alpha).$$



## Section C: Probability and Statistics

- 11** A coin is tossed repeatedly. The probability that a head appears is  $p$  and the probability that a tail appears is  $q = 1 - p$ .

- (i)**  $A$  and  $B$  play a game. The game ends if two successive heads appear, in which case  $A$  wins, or if two successive tails appear, in which case  $B$  wins.

Show that the probability that the game never ends is 0.

Given that the first toss is a head, show that the probability that  $A$  wins is  $\frac{p}{1 - pq}$ .

Find and simplify an expression for the probability that  $A$  wins.

- (ii)**  $A$  and  $B$  play another game. The game ends if three successive heads appear, in which case  $A$  wins, or if three successive tails appear, in which case  $B$  wins.

Show that

$$\Pr(A \text{ wins} \mid \text{the first toss is a head}) = p^2 + (q + pq) \Pr(A \text{ wins} \mid \text{the first toss is a tail})$$

and give a similar result for  $\Pr(A \text{ wins} \mid \text{the first toss is a tail})$ .

Show that

$$\Pr(A \text{ wins}) = \frac{p^2(1 - q^3)}{1 - (1 - p^2)(1 - q^2)}.$$

- (iii)**  $A$  and  $B$  play a third game. The game ends if  $a$  successive heads appear, in which case  $A$  wins, or if  $b$  successive tails appear, in which case  $B$  wins, where  $a$  and  $b$  are integers greater than 1.

Find the probability that  $A$  wins this game.

Verify that your result agrees with part **(i)** when  $a = b = 2$ .

- 12 The score shown on a biased  $n$ -sided die is represented by the random variable  $X$  which has distribution

$$\Pr(X = i) = \frac{1}{n} + \varepsilon_i \quad \text{for } i = 1, 2, \dots, n,$$

where not all the  $\varepsilon_i$  are equal to 0.

- (i) Find the probability that, when the die is rolled twice, the same score is shown on both rolls. Hence determine whether it is more likely for a fair die or a biased die to show the same score on two successive rolls.
- (ii) Use part (i) to prove that, for any set of  $n$  positive numbers  $x_i$  ( $i = 1, 2, \dots, n$ ),

$$\sum_{i=2}^n \sum_{j=1}^{i-1} x_i x_j \leq \frac{n-1}{2n} \left( \sum_{i=1}^n x_i \right)^2.$$

- (iii) Determine, with justification, whether it is more likely for a fair die or a biased die to show the same score on three successive rolls.